

**Table 8.6d**  
**Immunosuppression Use for Maintenance by Regimen**  
**Prior to Discharge, 1998 to 2007**  
**Recipients with Kidney-Pancreas Transplants**

	Transplant Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
<b>Functioning Graft at Discharge</b>	933	907	888	861	880	847	864	884	905	841
<b>With Immunosuppression Info</b>	933	907	886	861	880	844	850	879	891	826
CyA	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%
Tac	0.0%	0.7%	0.9%	1.4%	2.3%	1.1%	3.4%	0.8%	1.1%	1.1%
--- + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	1.1%	2.0%	1.0%	0.4%	0.0%
CyA + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	0.2%	0.7%	0.1%	0.4%
Tac + MMF/MPA	1.0%	0.6%	1.1%	0.6%	1.1%	4.0%	6.7%	14.8%	21.1%	24.1%
Tac + OtherAntimet	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.2%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.1%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	1.3%	2.8%	2.8%	3.1%	2.3%
Tac + Siro/Evero	0.0%	0.0%	0.3%	1.6%	4.7%	3.6%	3.8%	6.1%	2.2%	1.2%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	2.4%	4.5%	0.2%	0.2%	0.2%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.3%	0.2%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	1.5%	1.8%	2.1%	0.9%	1.8%	4.6%	4.6%
--- + Steroids	0.5%	2.2%	1.1%	0.1%	0.0%	1.7%	0.8%	2.2%	0.0%	0.1%
CyA + Steroids	0.5%	0.3%	0.5%	0.3%	0.1%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac + Steroids	2.4%	2.4%	1.7%	2.3%	1.8%	2.5%	1.5%	2.5%	3.0%	1.5%
--- + MMF/MPA + Steroids	0.9%	1.0%	0.1%	0.8%	0.5%	0.8%	1.3%	0.5%	0.7%	0.7%
CyA + MMF/MPA + Steroids	24.4%	19.4%	12.2%	8.5%	6.9%	5.0%	1.4%	1.8%	1.6%	1.2%
Tac + MMF/MPA + Steroids	57.4%	65.4%	64.6%	61.7%	66.1%	62.4%	64.7%	59.8%	53.6%	59.0%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%
CyA + OtherAntimet + Steroids	0.9%	0.1%	0.3%	0.1%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%
Tac + OtherAntimet + Steroids	1.7%	1.1%	1.0%	0.1%	0.5%	0.7%	0.2%	0.0%	0.6%	0.4%
--- + Siro/Evero + Steroids	0.1%	0.0%	0.0%	0.0%	0.2%	0.2%	0.4%	0.0%	0.0%	0.1%
CyA + Siro/Evero + Steroids	0.1%	0.0%	2.6%	1.6%	0.6%	1.8%	0.9%	0.9%	0.7%	0.4%
Tac + Siro/Evero + Steroids	0.0%	0.7%	6.8%	11.0%	7.3%	4.9%	2.2%	2.5%	4.4%	1.3%
Other Regimen	10.0%	6.2%	6.7%	8.1%	5.9%	4.1%	1.8%	1.0%	2.0%	1.3%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.